



Perinatal Depression

Progressive Case Conference

Trainee Guide

Contributors

Lucy A. Hutner, MD, Co-Founder and Chief Medical Officer, Phoebe
Sarah M. Nagle-Yang, MD, University of Colorado School of Medicine

Pre-Assessment Learning

Before you attend the classroom didactics on this module, please review some basic concepts of perinatal depression by viewing the following resources:

Epidemiology, risk factors, and etiology videos

[Perinatal Depression: Epidemiology](#)

[Perinatal Depression: Risk Factors](#)

[Perinatal Depression: Pathophysiology](#)

Screening and diagnosis videos

[Screening and Diagnosis 1](#)

[Screening and Diagnosis 2](#)

[Screening and Diagnosis 3](#)

[Screening and Diagnosis 4](#)

[Screening and Diagnosis 5](#)

Impact of depression in pregnancy

[Perinatal Depression: Impact](#)

Pre-Assessment Learning Objectives

After reviewing these resources, participants will be able to:

Describe the prevalence of depression in the antenatal and postpartum periods

Select appropriate screening methods for depression in the perinatal population

Overview

Depression in the perinatal period is a widespread but treatable psychiatric disorder. This module will focus on several facets of depression in both the antenatal and postpartum period. This module examines the epidemiology, risk factors, and etiology of perinatal depression, followed by a discussion of diagnosis and screening issues. This module also discusses the impact of depression on pregnancy, including obstetric and psychiatric outcomes. Last, this module focuses on the postpartum period as a particularly vulnerable time for the development of depression and how the risks might be mitigated.

Session

Pre-assessment Learning (prior to didactic session)

Group activity with videos: 30 minutes

Applying knowledge to a new case: 10 minutes

Session Learning Objectives

1. Describe the biological, psychological, and social risk factors for the development of depression in the perinatal period, identifying and comparing possible etiologies
2. Discuss the differential diagnoses of antenatal and postpartum depression
3. Discuss and explain the obstetric and psychiatric impact of depression in the perinatal period, including the potential for increased vulnerability during the postpartum period
4. Discuss a potential treatment approach for perinatal depression, encompassing biological, psychological, and social approaches

Case Presentation Part 1a: Guided Case Presentation

MR is a 27 year old G3P1 woman with a history of major depressive disorder and childhood trauma, who is currently 33 weeks pregnant, referred to you by her obstetrician after receiving an [EPDS](#) score of 22 during a recent prenatal visit. Current symptoms include 6 weeks of worsening mood, irritability, low motivation, poor appetite, poor concentration, restless sleep pattern and increased fatigue. She feels “incapacitated” by everyday life and feels “dark, like living is painful.” She endorses feelings of worthlessness and passive death wish. She denies active suicidal ideation, intent or plan.

She has no past suicide attempts or hospitalizations. Her history is significant for mild depressive episodes “on and off” since adolescence, often occurring in the days prior to menses and during major life transitions such as her first year of college and the transition to parenthood with her first child. Her most severe depressive episode occurred after the birth of her daughter 9 years ago. Despite being treated with Zoloft at that time with good effect, she has been anxious during this pregnancy about experiencing another severe episode and has felt “guilty about whether I’m even a good mother.”

Family history is notable for bipolar disorder in her mother. Her maternal grandmother also required psychiatric admissions following the birth of two of her four children.

Patient reports a harsh parenting style by her father, and physical abuse by both parents. She is currently estranged from both parents. She also was sexually abused by a male relative from age 10-13. She reports often “feeling alone” throughout her life. She currently lives with her fiancé, who is supportive. She is employed at a computer store as a tech specialist. She has been unable to work due to her depressive symptoms and has exhausted her short-term disability, leading to worry she could lose her job. Her fiancé has been picking up extra shifts at work to compensate financially, which has made her feel more distant from him and more socially isolated given that he is her primary social support.

Observe how the clinician discussed the possibility of the diagnosis of antenatal depression and the results of her EPDS score. [Clinical Case 1](#)

Case Presentation Part 1b: Guided Case Questions

1. What is her diagnosis?
2. What risk factors does this patient have for the development of perinatal depression?
3. What evidence supports an etiological role for the development of perinatal depression?

4. How would you characterize her suicide and infanticide risk?

Case Presentation Part 2: Guided Case Questions

1. What are the obstetric and psychiatric outcomes that the patient is at risk for, given her depression?
2. How can the psychiatrist counsel the patient to reduce the chance of her depression worsening in the postpartum period?

Watch how one clinician addressed the diagnosis of perinatal depression and the impact of depression on pregnancy and treatment planning with the patient. [Clinical Case 2](#)

Part 3a: Applying Knowledge to a New Case

RT is a 36-year-old woman, G1P1, currently five weeks postpartum, who presents to her first outpatient psychiatry appointment with depressed mood. She describes low mood with hopelessness, insomnia, irritability, and negative ruminations. She has difficulty initiating sleep (even when her newborn is sleeping) and experiences anxious ruminations about the baby's health. She had previously been stable on a moderate dose of fluoxetine, however at the advice of her family physician she had tapered and discontinued fluoxetine when first trying to conceive this pregnancy. She has a history of SSRI use in the past, including trials of sertraline (felt like a "robot"), citalopram, and fluoxetine. There is no history of suicide attempts or psychiatric hospitalizations. Family history is notable for mother with severe depression and suicide attempt. Patient works as an attorney and is married to a medical resident. Stressors include ventricular septal defect in the neonate, which was repaired after birth. Her older brother died of a medical illness when the patient was eleven years old. She is exclusively breastfeeding her infant.

Part 3b: New Case Questions

1. What specific risk factors may be contributing to this patient's depression in the postpartum period?
2. What are some risks associated with postpartum depression?

3. What options might the psychiatrist consider in developing a treatment plan for RT?



References

- Bloch M, Schmidt PJ, Danaceau M, Murphy J, Nieman L, Rubinow DR. Effects of gonadal steroids in women with a history of postpartum depression. *Am J Psychiatry* 2000;157:924-930
- Brown JV, Wilson CA, Ayre K, Robertson L, South E, Molyneaux E, Trevillion K, Howard LM, Khalifeh H. Antidepressant treatment for postnatal depression. *Cochrane Database of Systematic Reviews* 2021, Issue 2. Art. No.: CD013560. DOI: 10.1002/14651858.CD013560.pub2. Accessed 20 March 2021.
- Byatt N, Masters GA, Bergman AL, Moore Simas TA. [Screening for Mental Health and Substance Use Disorders in Obstetric Settings](#). *Curr Psychiatry Rep.* 2020 Sep 16;22(11):62. Review.
- Committee on Obstetric Practice. "The American College of Obstetricians and Gynecologists Committee Opinion no. 630. Screening for perinatal depression." *Obstetrics and gynecology* 125.5 (2015): 1268.
- Couto TCE, Brancaglioni MYM, Alvim-Soares A, et al. Postpartum depression: a systematic review of the genetics involved. *World J Psychiatry* 2015;5:103-111
- Cox JL, Holden JM, Sagovsky R. Detection of postnatal depression: development of the 10-item Edinburgh Postnatal Depression Scale. *Br J Psychiatry* 1987;150:782-786
- Deligiannidis KM, Freeman MP. Complementary and alternative medicine therapies for perinatal depression. *Best Pract Res Clin Obstet Gynaecol* 2014;28:85-95
- Fairbrother N, Corbyn B, Thordarson DS, Ma A, Surm D. [Screening for perinatal anxiety disorders: Room to grow](#). *J Affect Disord.* 2019 Mar 8; 250:363-370.
- Gaynes BN, Gavin N, Meltzer-Brody S, et al. Perinatal depression: prevalence, screening accuracy, and screening outcomes. *Evid Rep Technol Assess (Summ)*2005:1-8
- Howard LM, Molyneaux E, Dennis C-L, Rochat T, Stein A, Milgrom J. Non-psychotic mental disorders in the perinatal period. *Lancet* 2014;384:1775-1788
- Milgrom, Jeannette, and Alan W. Gemmill. "Screening for perinatal depression." *Best Practice & Research Clinical Obstetrics & Gynaecology* 28.1 (2014): 13-23.
- Pearlstein, Teri. "Depression during pregnancy." *Best Practice & Research Clinical Obstetrics & Gynaecology* 29.5 (2015): 754-764.
- Siu AL; US Preventive Services Task Force (USPSTF), Bibbins-Domingo K, Grossman DC, Baumann LC, Davidson KW, Ebell M, García FA, Gillman M, Herzstein J, Kemper AR, Krist AH, Kurth AE, Owens DK, Phillips WR, Phipps MG, Pignone MP. Screening for Depression in Adults: US Preventive Services Task Force Recommendation Statement. *JAMA.* 2016 Jan 26;315(4):380-7. doi: 10.1001/jama.2015.18392. PMID: 26813211.
- Stewart, Donna E. "Depression during pregnancy." *New England Journal of Medicine* 365.17 (2011): 1605-1611.
- Stewart, Donna E., and Simone Vigod. "Postpartum depression." *New England Journal of Medicine* 375.22 (2016): 2177-2186.
- Wisner KL, Moses-Kolko EL, Sit DKY. Postpartum depression: a disorder in search of a definition. *Arch Womens Ment Health* 2010;13:37-40